

General Information

- Use to absorb moisture from broken skin around the stoma. This allows the skin barrier to adhere so the skin can heal
- Apply powder after the skin has been washed and thoroughly dried

Using Adapt stoma powder

1. Wash and dry your skin thoroughly.
2. Apply a small amount of Adapt stoma powder to the irritated *open* skin area (Figure 1). Reapply if needed to coat the irritated area.
3. Gently brush away excess powder using your hand or a soft tissue. The powder should stick only to the raw area and not be left on dry intact skin.
4. Apply your ostomy pouching system as usual.

Broken or irritated skin around your stoma is usually caused by one of these problems:*

- Leakage of discharge onto the skin
- Improperly sized skin barrier opening (too large, too small, wrong shape)
- Sensitivity to one of the products used on the skin
- Removing the pouch too frequently, which can tear the upper layer of skin
- Wounds next to the stoma resulting from disease or surgery

Frequently Asked Questions

Q: Should I use a barrier film prep or gel to seal the powder before applying my pouch?

A: Skin barrier wipes such as Skin Protective Wipes are **optional**. Most contain alcohol, which will cause a stinging sensation when applied to raw skin areas. Skin barrier wipes are not necessary to use over Adapt stoma powder, but your nurse may recommend them.

Q: Is there medication in Adapt stoma powder?

A: No, it simply absorbs excess moisture, allowing your pouch to stay in place on your skin. The skin usually heals when protected from stoma discharge.

Q: Can I use other powder such as talcum?

A: No, the use of cosmetic type powders will usually interfere with the seal on your pouch, causing leakage and further irritation.

Q: When should I stop using Adapt stoma powder?

A: Stop using the powder when the skin has healed and is no longer moist to the touch. Adapt stoma powder is not designed to be used for the prevention of skin irritation. Stop using the powder and contact your healthcare provider if your skin has not improved or is getting worse.

**Broken or irritated skin near the stoma should be evaluated by a healthcare professional who will help determine the cause of the skin problem and how to prevent further injury.*

For product questions, sampling needs, or detailed clinical questions concerning our products in the US call **1.888.740.8999**. In Canada call **1.800.263.7400**.




Figure 1



Adapt Stoma Powder
Stock No 7906



Hollister Incorporated
2000 Hollister Drive
Libertyville, Illinois 60048
1.800.323.4060

www.hollister.com

Hollister Limited
95 Mary Street
Aurora, Ontario L4G 1G3
1.800.263.7400