

Cat. No. 260100 / 260100B NDC 054365-400-07  
 Cat. No. 260103 / 260103B NDC 054365-400-08

**ChloroPrep® Single Swabstick/ChloroPrep® Triple Swabstick**

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
 Patient Preoperative Skin Preparation

**Non-sterile Solution**  
**Applicator is non-sterile**

1.75 ml APPLICATOR / 5.25 ml APPLICATOR

Not made with natural rubber latex

Do Not Reuse

Professional Use Only  
 External Use Only

**Drug Facts**

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

**Use** for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**

For external use only. Flammable, keep away from fire or flame.

- do not use with electrocautery procedures
- Do not use**
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**ChloroPrep® Single Swabstick/Triple Swabstick**

Chlorhexidine gluconate 2% w/v and isopropyl alcohol 70% v/v  
 Patient Preoperative Skin Preparation

**WARNING. FLAMMABLE.**  
**KEEP AWAY FROM FIRE OR FLAME.**

**Single Swabstick**

Cat No. 260100

Single Use

**Triple Swabstick**

Cat No. 260103

Single Use

**ChloroPrep® Single Swabstick/ChloroPrep® Triple Swabstick**

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

**Drug Facts** (continued)

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area using
  - one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm<sup>2</sup>)
  - three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm<sup>2</sup>)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds (when using the triple swabstick applicators, use each swabstick sequentially within the 30 seconds). Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes (when using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredients** USP purified water

**Questions?** ■ [www.chloroprep.com](http://www.chloroprep.com) ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

